

**Maryland Biological Stream Survey
2000-2004**

**Volume 8:
County Results**

Prepared for

Maryland Department of Natural Resources
Monitoring and Non-Tidal Assessment Division
Tawes State Office Building
580 Taylor Avenue
Annapolis, Maryland 21401

Prepared by:

Paul F. Kazyak
Maryland Department of Natural Resources
Tawes State Office Building
580 Taylor Avenue
Annapolis, Maryland 21401
and
Allison Brindley
Mark T. Southerland
Versar, Inc.
9200 Rumsey Road
Columbia, Maryland 21045

July 2005

FOREWORD

This report volume, 2000-2004 Maryland Biological Stream Survey Volume 8: County Results, was prepared by staff from the Maryland Department of Natural Resources' Monitoring and Non-Tidal Assessment Division. It was supported in part by Maryland's Power Plant Research Program (PPRP Contract No. K00B020019 to Versar, Inc.).

ACKNOWLEDGMENTS

All aspects of the Maryland Biological Stream Survey have been a cooperative effort among Maryland DNR, several academic institutions, and consulting firms, as listed in the Acknowledgments sections of Volumes 1-5 of this report. For this volume, the authors would also like to acknowledge the contributions of the following individuals who contributed data analyses, graphical support, editing, or other talents specific to this volume:

Versar	DNR
Lori Erb	Scott Stranko
Mark Southerland	Martin Hurd
Ginny Rogers	Jay Kilian
Ed Weber	Ann Schenk
Allison Brindley	Daniel Boward
Sherian George	Brian Moody
Gail Lucas	

We would also like to thank Keith Van Ness, Bill Stack, Chris Victoria, Angela Morales, Shannon Moore, Tim Larney, Lynn Davidson, and Gwenda Brewer for serving as external peer reviewers for all or part of this volume.

ABSTRACT

One of the most important potential uses of the Maryland Biological Stream Survey (MBSS) is at the local level, by citizens and county governments. Recognizing this importance, this volume assessing 2000-2004 MBSS data was prepared to highlight survey results in a form most useable by local governments. Among all counties in Maryland, Queens Anne's County had the highest Combined Biotic Index (CBI) score (3.49), with an overall stream health rating of Fair. Other counties in the top five included Charles, Carroll, St. Mary's, and Talbot. In contrast, Baltimore City had the lowest average CBI (2.18), with an overall stream integrity rating of Poor. The next lowest counties were Somerset, Dorchester, Anne Arundel, and Washington. Charles and Prince George's counties shared the highest rated watershed for freshwater biodiversity (Zekiah Swamp), while Frederick County's Catoctin Creek watershed was the lowest ranked watershed for freshwater biodiversity in any county. Counties with low dissolved oxygen (DO) levels included: Somerset (38% of stream miles), Dorchester (26%), Caroline (26%), Baltimore City, and Talbot (23%). Caroline County had the highest mean nitrate-nitrogen levels (5.51 mg/l), followed by Dorchester County (4.59 mg/l); the lowest county mean was observed in Somerset County (0.40 mg/l). There was a strong relationship between mean nitrate-nitrogen levels for each county and the percentage of that county harvested as cropland. Allegany (0.010 mg/l) and Garrett Counties (0.013 mg/l) had the lowest mean levels of Total

Phosphorous, while Worcester (0.125 mg/l) and Kent (0.117 mg/l) had the highest mean levels. The total number of chickens in a county was significantly related to the mean Total Phosphorous concentration in that county. Allegheny and Garrett Counties had the highest mean Physical Habitat Index score (77 on a 100 point scale), while Baltimore City (50/100) and Cecil County (63/100) had the lowest mean habitat values. Urban counties had poorer trash rating scores than agricultural counties. Baltimore City received the lowest mean rating (5 on a 20 point scale), followed by Prince George's, Anne Arundel, and Baltimore County. The least amount of human refuse on average along or in streams was found in Garrett (18/20) and Queen Annes Counties (17/20). Among all counties, Washington had the highest estimate of stream miles with no riparian buffer (20%), followed by Wicomico (19%), Cecil (11%) and Frederick (11%). The counties with the highest occurrence of riparian buffer breaks were Baltimore City (44%), Wicomico (19%), Worcester (15%), and Baltimore County (15%). From these analyses, it is clear that buffer breaks are a potentially important limitation to the effectiveness of riparian buffers in Maryland. The information contained here should help policymakers, planners, and others justify and guide natural resource conservation and restoration efforts in Maryland. In addition, the findings presented here may help guide other restoration efforts into the most needed areas, especially those associated with the Chesapeake Bay.

TABLE OF CONTENTS

	Page
FOREWORD	iii
ACKNOWLEDGMENTS	v
ABSTRACT.....	vii
 8.1 INTRODUCTION	 8-1
8.2 COUNTY COMPARISONS	8-2
8.2.1 Biotic Integrity	8-2
8.2.2 Chemical Conditions	8-2
8.2.2.1 Dissolved Oxygen.....	8-2
8.2.2.2 Nitrogen	8-3
8.2.2.3 Phosphorous.....	8-3
8.2.2.4 pH and Acid Neutralizing Capacity (ANC).....	8-6
8.2.3 Physical Conditions.....	8-6
8.2.3.1 Overall Conditions.....	8-6
8.2.3.2 Trash	8-8
8.2.3.3 Channelization	8-8
8.2.3.4 Bank Erosion	8-9
8.2.4 Riparian Conditions	8-9
8.2.5 County Stressors.....	8-9
 8.3 ALLEGANY COUNTY	 8-13
8.3.1 Ecological Health	8-13
8.3.2 Physical Habitat	8-13
8.3.2.1 Overall Condition	8-13
8.3.2.2 Trash	8-13
8.3.2.3 Channelization	8-13
8.3.2.4 Inadequate Riparian Buffer.....	8-13
8.3.2.5 Eroded Banks/Bedload Movement	8-14
8.3.3 Key Nutrients	8-14
8.3.3.1 Nitrate-Nitrogen.....	8-14
8.3.3.2 Total Phosphorus	8-14
8.3.4 Stream and River Biodiversity	8-14
8.3.5 Stressors	8-14
 8.4 ANNE ARUNDEL COUNTY	 8-27
8.4.1 Ecological Health	8-27
8.4.2 Physical Habitat	8-27
8.4.2.1 Overall Condition	8-27
8.4.2.2 Trash	8-27
8.4.2.3 Channelization	8-27
8.4.2.4 Inadequate Riparian Buffer.....	8-27
8.4.2.5 Eroded Banks/Bedload Movement	8-27
8.4.3 Key Nutrients	8-28
8.4.3.1 Nitrate-Nitrogen.....	8-28
8.4.3.2 Total Phosphorus	8-28
8.4.4 Stream and River Biodiversity	8-28
8.4.5 Stressors	8-28
 8.5 BALTIMORE COUNTY	 8-41
8.5.1 Ecological Health	8-41
8.5.2 Physical Habitat	8-41
8.5.2.1 Overall Condition	8-41
8.5.2.2 Trash	8-42
8.5.2.3 Channelization	8-42
8.5.2.4 Inadequate Riparian Buffer.....	8-42

TABLE OF CONTENTS (CONTINUED)

	Page
8.5.2.5 Eroded Banks/Bedload Movement	8-42
8.5.3 Key Nutrients	8-42
8.5.3.1 Nitrate-Nitrogen.....	8-42
8.5.3.2 Total Phosphorus	8-42
8.5.4 Stream and River Biodiversity	8-42
8.5.5 Stressors	8-43
8.6 BALTIMORE CITY	8-57
8.6.1 Ecological Health	8-57
8.6.2 Physical Habitat	8-57
8.6.2.1 Overall Condition	8-57
8.6.2.2 Trash	8-57
8.6.2.3 Channelization	8-57
8.6.2.4 Inadequate Riparian Buffer.....	8-57
8.6.2.5 Eroded Banks/Bedload Movement	8-57
8.6.3 Key Nutrients	8-58
8.6.3.1 Nitrate-Nitrogen.....	8-58
8.6.3.2 Total Phosphorus	8-58
8.6.4 Stream and River Biodiversity	8-58
8.6.5 Stressors	8-58
8.7 CALVERT COUNTY	8-71
8.7.1 Ecological Health	8-71
8.7.2 Physical Habitat	8-71
8.7.2.1 Overall Condition	8-71
8.7.2.2 Trash	8-71
8.7.2.3 Channelization	8-71
8.7.2.4 Inadequate Riparian Buffer.....	8-71
8.7.2.5 Eroded Banks/Bedload Movement	8-71
8.7.3 Key Nutrients	8-72
8.7.3.1 Nitrate-Nitrogen.....	8-72
8.7.3.2 Total Phosphorus	8-72
8.7.4 Stream and River Biodiversity	8-72
8.7.5 Stressors	8-72
8.8 CAROLINE COUNTY	8-85
8.8.1 Ecological Health	8-85
8.8.2 Physical Habitat	8-85
8.8.2.1 Overall Condition	8-85
8.8.2.2 Trash	8-85
8.8.2.3 Channelization	8-85
8.8.2.4 Inadequate Riparian Buffer.....	8-85
8.8.2.5 Eroded Banks/Bedload Movement	8-86
8.8.3 Key Nutrients	8-86
8.8.3.1 Nitrate-Nitrogen.....	8-86
8.8.3.2 Total Phosphorus	8-86
8.8.4 Stream and River Biodiversity	8-86
8.8.5 Stressors	8-86
8.9 CARROLL COUNTY	8-101
8.9.1 Ecological Health	8-101
8.9.2 Physical Habitat	8-101
8.9.2.1 Overall Condition	8-101
8.9.2.2 Trash	8-101
8.9.2.3 Channelization	8-102

TABLE OF CONTENTS (CONTINUED)

	Page
8.9.2.4 Inadequate Riparian Buffer.....	8-102
8.9.2.5 Eroded Banks/Bedload Movement	8-102
8.9.3 Key Nutrients	8-102
8.9.3.1 Nitrate-Nitrogen.....	8-102
8.9.3.2 Total Phosphorus	8-102
8.9.4 Stream and River Biodiversity	8-102
8.9.5 Stressors	8-103
8.10 CECIL COUNTY	8-117
8.10.1 Ecological Health	8-117
8.10.2 Physical Habitat	8-117
8.10.2.1 Overall Condition	8-117
8.10.2.2 Trash	8-117
8.10.2.3 Channelization	8-117
8.10.2.4 Inadequate Riparian Buffer.....	8-118
8.10.2.5 Eroded Banks/Bedload Movement	8-118
8.10.3 Key Nutrients	8-118
8.10.3.1 Nitrate-Nitrogen.....	8-118
8.10.3.2 Total Phosphorus	8-118
8.10.4 Stream and River Biodiversity	8-118
8.10.5 Stressors	8-119
8.11 CHARLES COUNTY.....	8-133
8.11.1 Ecological Health	8-133
8.11.2 Physical Habitat	8-133
8.11.2.1 Overall Condition	8-133
8.11.2.2 Trash	8-133
8.11.2.3 Channelization	8-133
8.11.2.4 Inadequate Riparian Buffer.....	8-133
8.11.2.5 Eroded Banks/Bedload Movement	8-133
8.11.3 Key Nutrients	8-134
8.11.3.1 Nitrate-Nitrogen.....	8-134
8.11.3.2 Total Phosphorus	8-134
8.11.4 Stream and River Biodiversity	8-134
8.11.5 Stressors	8-134
8.12 DORCHESTER COUNTY	8-147
8.12.1 Ecological Health	8-147
8.12.2 Physical Habitat	8-147
8.12.2.1 Overall Condition	8-147
8.12.2.2 Trash	8-147
8.12.2.3 Channelization	8-147
8.12.2.4 Inadequate Riparian Buffer.....	8-147
8.12.2.5 Eroded Banks/Bedload Movement	8-147
8.12.3 Key Nutrients	8-148
8.12.3.1 Nitrate-Nitrogen.....	8-148
8.12.3.2 Total Phosphorus	8-148
8.12.4 Stream and River Biodiversity	8-148
8.12.5 Stressors	8-148
8.13 FREDERICK COUNTY	8-161
8.13.1 Ecological Health	8-161
8.13.2 Physical Habitat	8-161
8.13.2.1 Overall Condition	8-161
8.13.2.2 Trash	8-161

TABLE OF CONTENTS (CONTINUED)

	Page
8.13.2.3 Channelization	8-162
8.13.2.4 Inadequate Riparian Buffer.....	8-162
8.13.2.5 Eroded Banks/Bedload Movement	8-162
8.13.3 Key Nutrients	8-162
8.13.3.1 Nitrate-Nitrogen.....	8-162
8.13.3.2 Total Phosphorus	8-162
8.13.4 Stream and River Biodiversity	8-162
8.13.5 Stressors	8-163
8.14 GARRETT COUNTY	8-177
8.14.1 Ecological Health	8-177
8.14.2 Physical Habitat	8-177
8.14.2.1 Overall Condition	8-177
8.14.2.2 Trash	8-177
8.14.2.3 Channelization	8-177
8.14.2.4 Inadequate Riparian Buffer.....	8-177
8.14.2.5 Eroded Banks/Bedload Movement	8-178
8.14.3 Key Nutrients	8-178
8.14.3.1 Nitrate-Nitrogen.....	8-178
8.14.3.2 Total Phosphorus	8-178
8.14.4 Stream and River Biodiversity	8-178
8.14.5 Stressors	8-179
8.15 HARFORD COUNTY	8-193
8.15.1 Ecological Health	8-193
8.15.2 Physical Habitat	8-193
8.15.2.1 Overall Condition	8-193
8.15.2.2 Trash	8-193
8.15.2.3 Channelization	8-193
8.15.2.4 Inadequate Riparian Buffer.....	8-193
8.15.2.5 Eroded Banks/Bedload Movement	8-193
8.15.3 Key Nutrients	8-194
8.15.3.1 Nitrate-Nitrogen.....	8-194
8.15.3.2 Total Phosphorus	8-194
8.15.4 Stream and River Biodiversity	8-194
8.15.5 Stressors	8-194
8.16 HOWARD COUNTY	8-209
8.16.1 Ecological Health	8-209
8.16.2 Physical Habitat	8-209
8.16.2.1 Overall Condition	8-209
8.16.2.2 Trash	8-209
8.16.2.3 Channelization	8-210
8.16.2.4 Inadequate Riparian Buffer.....	8-210
8.16.2.5 Eroded Banks/Bedload Movement	8-210
8.16.3 Key Nutrients	8-210
8.16.3.1 Nitrate-Nitrogen.....	8-210
8.16.3.2 Total Phosphorus	8-210
8.16.4 Stream and River Biodiversity	8-210
8.16.5 Stressors	8-211
8.17 KENT COUNTY	8-223
8.17.1 Ecological Health	8-223
8.17.2 Physical Habitat	8-223
8.17.2.1 Overall Condition	8-223

TABLE OF CONTENTS (CONTINUED)

	Page
8.17.2.2 Trash	8-223
8.17.2.3 Channelization	8-223
8.17.2.4 Inadequate Riparian Buffer.....	8-223
8.17.2.5 Eroded Banks/Bedload Movement	8-223
8.17.3 Key Nutrients	8-224
8.17.3.1 Nitrate-Nitrogen.....	8-224
8.17.3.2 Total Phosphorus	8-224
8.17.4 Stream and River Biodiversity	8-224
8.17.5 Stressors	8-224
8.18 MONTGOMERY COUNTY	8-237
8.18.1 Ecological Health	8-237
8.18.2 Physical Habitat	8-237
8.18.2.1 Overall Condition	8-237
8.18.2.2 Trash	8-237
8.18.2.3 Channelization	8-238
8.18.2.4 Inadequate Riparian Buffer.....	8-238
8.18.2.5 Eroded Banks/Bedload Movement	8-238
8.18.3 Key Nutrients	8-238
8.18.3.1 Nitrate-Nitrogen.....	8-238
8.18.3.2 Total Phosphorus	8-238
8.18.4 Stream and River Biodiversity	8-238
8.18.5 Stressors	8-239
8.19 PRINCE GEORGE’S COUNTY.....	8-253
8.19.1 Ecological Health	8-253
8.19.2 Physical Habitat	8-253
8.19.2.1 Overall Condition	8-253
8.19.2.2 Trash	8-253
8.19.2.3 Channelization	8-253
8.19.2.4 Inadequate Riparian Buffer.....	8-253
8.19.2.5 Eroded Banks/Bedload Movement	8-253
8.19.3 Key Nutrients	8-254
8.19.3.1 Nitrate-Nitrogen.....	8-254
8.19.3.2 Total Phosphorus	8-254
8.19.4 Stream and River Biodiversity	8-254
8.19.5 Stressors	8-254
8.20 QUEEN ANNE’S COUNTY	8-267
8.20.1 Ecological Health	8-267
8.20.2 Physical Habitat	8-267
8.20.2.1 Overall Condition	8-267
8.20.2.2 Trash	8-267
8.20.2.3 Channelization	8-267
8.20.2.4 Inadequate Riparian Buffer.....	8-267
8.20.2.5 Eroded Banks/Bedload Movement	8-267
8.20.3 Key Nutrients	8-268
8.20.3.1 Nitrate-Nitrogen.....	8-268
8.20.3.2 Total Phosphorus	8-268
8.20.4 Stream and River Biodiversity	8-268
8.20.5 Stressors	8-268
8.21 SOMERSET COUNTY.....	8-281
8.21.1 Ecological Health	8-281
8.21.2 Physical Habitat	8-281

TABLE OF CONTENTS (CONTINUED)

	Page
8.21.2.1 Overall Condition	8-281
8.21.2.2 Trash	8-281
8.21.2.3 Channelization	8-281
8.21.2.4 Inadequate Riparian Buffer.....	8-281
8.21.2.5 Eroded Banks/Bedload Movement	8-281
8.21.3 Key Nutrients	8-282
8.21.3.1 Nitrate-Nitrogen.....	8-282
8.21.3.2 Total Phosphorus	8-282
8.21.4 Stream and River Biodiversity	8-282
8.21.5 Stressors	8-282
8.22 ST. MARY’S COUNTY	8-295
8.22.1 Ecological Health	8-295
8.22.2 Physical Habitat	8-295
8.22.2.1 Overall Condition	8-295
8.22.2.2 Trash	8-295
8.22.2.3 Channelization	8-295
8.22.2.4 Inadequate Riparian Buffer.....	8-295
8.22.2.5 Eroded Banks/Bedload Movement	8-296
8.22.3 Key Nutrients	8-296
8.22.3.1 Nitrate-Nitrogen.....	8-296
8.22.3.2 Total Phosphorus	8-296
8.22.4 Stream and River Biodiversity	8-296
8.22.5 Stressors	8-296
8.23 TALBOT COUNTY	8-311
8.23.1 Ecological Health	8-311
8.23.2 Physical Habitat	8-311
8.23.2.1 Overall Condition	8-311
8.23.2.2 Trash	8-311
8.23.2.3 Channelization	8-311
8.23.2.4 Inadequate Riparian Buffer.....	8-311
8.23.2.5 Eroded Banks/Bedload Movement	8-311
8.23.3 Key Nutrients	8-311
8.23.3.1 Nitrate-Nitrogen.....	8-311
8.23.3.2 Total Phosphorus	8-312
8.23.4 Stream and River Biodiversity	8-312
8.23.5 Stressors	8-312
8.24 WASHINGTON COUNTY.....	8-325
8.24.1 Ecological Health	8-325
8.24.2 Physical Habitat	8-325
8.24.2.1 Overall Condition	8-325
8.24.2.2 Trash	8-325
8.24.2.3 Channelization	8-325
8.24.2.4 Inadequate Riparian Buffer.....	8-325
8.24.2.5 Eroded Banks/Bedload Movement	8-325
8.24.3 Key Nutrients	8-326
8.24.3.1 Nitrate-Nitrogen.....	8-326
8.24.3.2 Total Phosphorus	8-326
8.24.4 Stream and River Biodiversity	8-326
8.24.5 Stressors	8-326

TABLE OF CONTENTS (CONTINUED)

	Page
8.25 WICOMICO COUNTY	8-339
8.25.1 Ecological Health	8-339
8.25.2 Physical Habitat	8-339
8.25.2.1 Overall Condition	8-339
8.25.2.2 Trash	8-339
8.25.2.3 Channelization	8-339
8.25.2.4 Inadequate Riparian Buffer	8-339
8.25.2.5 Eroded Banks/Bedload Movement	8-340
8.25.3 Key Nutrients	8-340
8.25.3.1 Nitrate-Nitrogen	8-340
8.25.3.2 Total Phosphorus	8-340
8.25.4 Stream and River Biodiversity	8-340
8.25.5 Stressors	8-341
 8.26 WORCESTER COUNTY	 8-355
8.26.1 Ecological Health	8-355
8.26.2 Physical Habitat	8-355
8.26.2.1 Overall Condition	8-355
8.26.2.2 Trash	8-355
8.26.2.3 Channelization	8-355
8.26.2.4 Inadequate Riparian Buffer	8-355
8.26.2.5 Eroded Banks/Bedload Movement	8-356
8.26.3 Key Nutrients	8-356
8.26.3.1 Nitrate-Nitrogen	8-356
8.26.3.2 Total Phosphorus	8-356
8.26.4 Stream and River Biodiversity	8-356
8.26.5 Stressors	8-356
 8.27 REFERENCES	 8-371

LIST OF TABLES

Table No.	Page
8-1 Mean biological integrity scores for 4 th order and smaller Maryland streams, by county, based on results from the 2000-2004 MBSS.....	8-3
8-2 Mean values for key chemical constituents of 4th order and smaller Maryland streams, by county, based on results from the 2000-2004 MBSS	8-4
8-3 Mean values for key physical attributes of 4th order and smaller Maryland streams, by county, based on results from the 2000-2004 MBSS	8-7
8-4 Estimates of channelization types by county based on the 2000-2004 MBSS	8-8
8-5 MBSS sites sampled in Allegany County during 1994-2004, ranked by Combined Biotic Index Score	8-17
8-6 Stream Waders sites sampled in Allegany County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-18
8-7 MBSS sites sampled in Anne Arundel County during 1994-2004, ranked by Combined Biotic Index Score	8-31
8-8 Stream Waders sites sampled in Anne Arundel County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-32
8-9 MBSS sites sampled in Baltimore County during 1994-2004, ranked by Combined Biotic Index Score	8-46
8-10 Stream Waders sites sampled in Baltimore County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-47
8-11 MBSS sites sampled in Baltimore City during 1994-2004, ranked by Combined Biotic Index Score.....	8-61
8-12 Stream Waders sites sampled in Baltimore City during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-62
8-13 MBSS sites sampled in Calvert County during 1994-2004, ranked by Combined Biotic Index Score	8-75
8-14 Stream Waders sites sampled in Calvert County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-75
8-15 MBSS sites sampled in Caroline County during 1994-2004, ranked by Combined Biotic Index Score	8-90
8-16 Stream Waders sites sampled in Caroline County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-91
8-17 MBSS sites sampled in Carroll County during 1994-2004, ranked by Combined Biotic Index Score.....	8-106
8-18 Stream Waders sites sampled in Carroll County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-107
8-19 MBSS sites sampled in Cecil County during 1994-2004, ranked by Combined Biotic Index Score	8-122
8-20 Stream Waders sites sampled in Cecil County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-123
8-21 MBSS sites sampled in Charles County during 1994-2004, ranked by Combined Biotic Index Score.....	8-137
8-22 Stream Waders sites sampled in Charles County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-138

LIST OF TABLES (CONTINUED)

Table No.	Page
8-23 MBSS sites sampled in Dorchester County during 1994-2004, ranked by Combined Biotic Index Score	8-151
8-24 Stream Waders sites sampled in Dorchester County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-152
8-25 MBSS sites sampled in Frederick County during 1994-2004, ranked by Combined Biotic Index Score.....	8-166
8-26 Stream Waders sites sampled in Frederick County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-167
8-27 MBSS sites sampled in Garrett County during 1994-2004, ranked by Combined Biotic Index Score.....	8-182
8-28 Stream Waders sites sampled in Garrett County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-183
8-29 MBSS sites sampled in Harford County during 1994-2004, ranked by Combined Biotic Index Score	8-198
8-30 Stream Waders sites sampled in Harford County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-199
8-31 MBSS sites sampled in Howard County during 1994-2004, ranked by Combined Biotic Index Score	8-212
8-32 Stream Waders sites sampled in Howard County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-213
8-33 MBSS sites sampled in Kent County during 1994-2004, ranked by Combined Biotic Index Score	8-227
8-34 Stream Waders sites sampled in Kent County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-228
8-35 MBSS sites sampled in Montgomery County during 1994-2004, ranked by Combined Biotic Index Score	8-242
8-36 Stream Waders sites sampled in Montgomery County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-243
8-37 MBSS sites sampled in Prince George's County during 1994-2004, ranked by Combined Biotic Index Score	8-257
8-38 Stream Waders sites sampled in Prince George's County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-258
8-39 MBSS sites sampled in Queen Anne's County during 1994-2004, ranked by Combined Biotic Index Score	8-271
8-40 Stream Waders sites sampled in Queen Anne's County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-272
8-41 MBSS sites sampled in Somerset County during 1994-2004, ranked by Combined Biotic Index Score	8-285
8-42 Stream Waders sites sampled in Somerset County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-286
8-43 MBSS sites sampled in St. Mary's County during 1994-2004, ranked by Combined Biotic Index Score	8-300
8-44 Stream Waders sites sampled in St. Mary's County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity	8-301
8-xviii	

LIST OF TABLES (CONTINUED)

Table No.	Page
8-45	MBSS sites sampled in Talbot County during 1994-2004, ranked by Combined Biotic Index Score8-315
8-46	Stream Waders sites sampled in Talbot County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity8-316
8-47	MBSS sites sampled in Washington County during 1994-2004, ranked by Combined Biotic Index Score.....8-329
8-48	Stream Waders sites sampled in Washington County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity8-330
8-49	MBSS sites sampled in Wicomico County during 1994-2004, ranked by Combined Biotic Index Score8-344
8-50	Stream Waders sites sampled in Wicomico County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity8-345
8-51	MBSS sites sampled in Worcester County during 1994-2004, ranked by Combined Biotic Index Score8-360
8-52	Stream Waders sites sampled in Worcester County during 2000-2004, ranked by Family-level Benthic Index of Biotic Integrity8-361

LIST OF FIGURES

Figure No.		Page
8-1	Dissolved organic carbon levels vs. dissolved oxygen from the 2000-2004 MBSS.....	8-5
8-2	Mean nitrate-nitrogen levels from the 2000-2004 MBSS, by county, and the percentage of that county harvested as cropland	8-5
8-3	Mean Total Phosphorous levels from the 2000-2004 MBSS, by county, and the number of chickens raised in the county	8-5
8-4	Relative severity of stressors affecting biota in Maryland streams, based on 2000-2004 MBSS data.....	8-11
8-5	Stressors by county, in percent of total stream miles for that county, based on the 2000-2004 MBSS	8-12
ALLEGANY COUNTY		
8-6	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Allegany County streams sampled by the MBSS during 1995-97 and 2000-2004	8-16
8-7	Physical Habitat Index pie chart and map of stream habitat quality for Allegany County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-19
8-8	Pie chart and map of trash rating for Allegany County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-20
8-9	Map of channelized sites, by type, for Allegany County streams sampled by the MBSS during 2000-2004.....	8-21
8-10	Map of sites with inadequate riparian buffers and buffer breaks for Allegany County streams sampled by the MBSS during 2000-2004.....	8-22
8-11	Pie charts and map of sites with eroded banks and instream bar formation for Allegany County streams sampled by the MBSS during 2000-2004.....	8-23
8-12	Pie chart and map of nitrate-nitrogen values for Allegany County streams sampled by the MBSS during 1995-97 and 2000-2004	8-24
8-13	Pie chart and map of total phosphorus values for Allegany County streams sampled by the MBSS during 2000-2004.....	8-25
8-14	Aquatic Heritage Biodiversity Ranking map for Allegany County, by watershed.....	8-26
ANNE ARUNDEL COUNTY		
8-15	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Anne Arundel County streams sampled by the MBSS during 1995-97 and 2000-2004	8-30
8-16	Physical Habitat Index pie chart and map of stream habitat quality for Anne Arundel County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-33
8-17	Pie chart and map of trash rating for Anne Arundel County streams sampled by the MBSS during 1995-97 and 2000-2004	8-34
8-18	Map of channelized sites, by type, for Anne Arundel County streams sampled by the MBSS during 2000-2004	8-35
8-19	Map of sites with inadequate riparian buffers and buffer breaks for Anne Arundel County streams sampled by the MBSS during 2000-2004.....	8-36
8-20	Pie charts and map of sites with eroded banks and instream bar formation for Anne Arundel County streams sampled by the MBSS during 2000-2004.....	8-37
8-21	Pie chart and map of nitrate-nitrogen values for Anne Arundel County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-38
8-22	Pie chart and map of total phosphorus values for Anne Arundel County streams sampled by the MBSS during 2000-2004.....	8-39
8-23	Aquatic Heritage Biodiversity Ranking map for Anne Arundel County, by watershed.....	8-40
BALTIMORE COUNTY		
8-24	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Baltimore County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-45
8-25	Physical Habitat Index pie chart and map of stream habitat quality for Baltimore County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-48
8-26	Pie chart and map of trash rating for Baltimore County streams sampled by the MBSS during 1995-97 and 2000-2004	8-49
8-xx		

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-27	Map of channelized sites, by type, for Baltimore County streams sampled by the MBSS during 2000-2004.....8-50
8-28	Map of sites with inadequate riparian buffers and buffer breaks for Baltimore County streams sampled by the MBSS during 2000-2004.....8-51
8-29	Pie charts and map of sites with eroded banks and instream bar formation for Baltimore County streams sampled by the MBSS during 2000-2004.....8-52
8-30	Pie chart and map of nitrate-nitrogen values for Baltimore County streams sampled by the MBSS during 1995-97 and 2000-20048-53
8-31	Pie chart and map of total phosphorus values for Baltimore County streams sampled by the MBSS during 2000-20048-54
8-32	Aquatic Heritage Biodiversity Ranking map for Baltimore County, by watershed.....8-55
BALTIMORE CITY	
8-33	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Baltimore City streams sampled by the MBSS during 1995-97 and 2000-20048-60
8-34	Physical Habitat Index pie chart and map of stream habitat quality for Baltimore City streams sampled by the MBSS during 1995-97 and 2000-20048-63
8-35	Pie chart and map of trash rating for Baltimore City streams sampled by the MBSS during 1995-97 and 2000-20048-64
8-36	Map of channelized sites, by type, for Baltimore City streams sampled by the MBSS during 2000-2004.....8-65
8-37	Map of sites with inadequate riparian buffers and buffer breaks for Baltimore City streams sampled by the MBSS during 2000-2004.....8-66
8-38	Pie charts and map of sites with eroded banks and instream bar formation for Baltimore City streams sampled by the MBSS during 2000-2004.....8-67
8-39	Pie chart and map of nitrate-nitrogen values for Baltimore City streams sampled by the MBSS during 1995-97 and 2000-2004.....8-68
8-40	Pie chart and map of total phosphorus values for Baltimore City streams sampled by the MBSS during 2000-20048-69
8-41	Aquatic Heritage Biodiversity Ranking map for Baltimore City, by watershed.....8-70
CALVERT COUNTY	
8-42	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Calvert County streams sampled by the MBSS during 1995-97 and 2000-20048-74
8-43	Physical Habitat Index pie chart and map of stream habitat quality for Calvert County streams sampled by the MBSS during 1995-97 and 2000-20048-76
8-44	Pie chart and map of trash rating for Calvert County streams sampled by the MBSS during 1995-97 and 2000-20048-77
8-45	Map of channelized sites, by type, for Calvert County streams sampled by the MBSS during 2000-20048-78
8-46	Map of sites with inadequate riparian buffers and buffer breaks for Calvert County streams sampled by the MBSS during 2000-2004.....8-79
8-47	Pie charts and map of sites with eroded banks and instream bar formation for Calvert County streams sampled by the MBSS during 2000-2004.....8-80
8-48	Pie chart and map of nitrate-nitrogen values for Calvert County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-81
8-49	Pie chart and map of total phosphorus values for Calvert County streams sampled by the MBSS during 2000-2004.....8-82
8-50	Aquatic Heritage Biodiversity Ranking map for Calvert County, by watershed.....8-83
CAROLINE COUNTY	
8-51	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Caroline County streams sampled by the MBSS during 1995-97 and 2000-20048-89
8-52	Physical Habitat Index pie chart and map of stream habitat quality for Caroline County streams sampled by the MBSS during 1995-97 and 2000-20048-92
8-53	Pie chart and map of trash rating for Caroline County streams sampled by the MBSS during 1995-97 and 2000-20048-93

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-54	Map of channelized sites, by type, for Caroline County streams sampled by the MBSS during 2000-20048-94
8-55	Map of sites with inadequate riparian buffers and buffer breaks for Caroline County streams sampled by the MBSS during 2000-20048-95
8-56	Pie charts and map of sites with eroded banks and instream bar formation for Caroline County streams sampled by the MBSS during 2000-20048-96
8-57	Pie chart and map of nitrate-nitrogen values for Caroline County streams sampled by the MBSS during 1995-97 and 2000-20048-97
8-58	Pie chart and map of total phosphorus values for Caroline County streams sampled by the MBSS during 2000-20048-98
8-59	Aquatic Heritage Biodiversity Ranking map for Caroline County, by watershed8-99
CARROLL COUNTY	
8-60	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Carroll County streams sampled by the MBSS during 1995-97 and 2000-20048-105
8-61	Physical Habitat Index pie chart and map of stream habitat quality for Carroll County streams sampled by the MBSS during 1995-97 and 2000-20048-108
8-62	Pie chart and map of trash rating for Carroll County streams sampled by the MBSS during 1995-97 and 2000-20048-109
8-63	Map of channelized sites, by type, for Carroll County streams sampled by the MBSS during 2000-20048-110
8-64	Map of sites with inadequate riparian buffers and buffer breaks for Carroll County streams sampled by the MBSS during 2000-20048-111
8-65	Pie charts and map of sites with eroded banks and instream bar formation for Carroll County streams sampled by the MBSS during 2000-20048-112
8-66	Pie chart and map of nitrate-nitrogen values for Carroll County streams sampled by the MBSS during 1995-97 and 2000-20048-113
8-67	Pie chart and map of total phosphorus values for Carroll County streams sampled by the MBSS during 2000-20048-114
8-68	Aquatic Heritage Biodiversity Ranking map for Carroll County, by watershed8-115
CECIL COUNTY	
8-69	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Cecil County streams sampled by the MBSS during 1995-97 and 2000-20048-121
8-70	Physical Habitat Index pie chart and map of stream habitat quality for Cecil County streams sampled by the MBSS during 1995-97 and 2000-20048-124
8-71	Pie chart and map of trash rating for Cecil County streams sampled by the MBSS during 1995-97 and 2000-20048-125
8-72	Map of channelized sites, by type, for Cecil County streams sampled by the MBSS during 2000-20048-126
8-73	Map of sites with inadequate riparian buffers and buffer breaks for Cecil County streams sampled by the MBSS during 2000-20048-127
8-74	Pie charts and map of sites with eroded banks and instream bar formation for Cecil County streams sampled by the MBSS during 2000-20048-128
8-75	Pie chart and map of nitrate-nitrogen values for Cecil County streams sampled by the MBSS during 1995-97 and 2000-20048-129
8-76	Pie chart and map of total phosphorus values for Cecil County streams sampled by the MBSS during 2000-20048-130
8-77	Aquatic Heritage Biodiversity Ranking map for Cecil County, by watershed8-131
CHARLES COUNTY	
8-78	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Charles County streams sampled by the MBSS during 1995-97 and 2000-20048-136
8-79	Physical Habitat Index pie chart and map of stream habitat quality for Charles County streams sampled by the MBSS during 1995-97 and 2000-20048-139
8-80	Pie chart and map of trash rating for Charles County streams sampled by the MBSS during 1995-97 and 2000-20048-140
8-81	Map of channelized sites, by type, for Charles County streams sampled by the MBSS during 2000-20048-141

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-82	Map of sites with inadequate riparian buffers and buffer breaks for Charles County streams sampled by the MBSS during 2000-2004.....8-142
8-83	Pie charts and map of sites with eroded banks and instream bar formation for Charles County streams sampled by the MBSS during 2000-2004.....8-143
8-84	Pie chart and map of nitrate-nitrogen values for Charles County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-144
8-85	Pie chart and map of total phosphorus values for Charles County streams sampled by the MBSS during 2000-2004.....8-145
8-86	Aquatic Heritage Biodiversity Ranking map for Charles County, by watershed8-146
DORCHESTER COUNTY	
8-87	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Dorchester County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-150
8-88	Physical Habitat Index pie chart and map of stream habitat quality for Dorchester County streams sampled by the MBSS during 1995-97 and 2000-20048-153
8-89	Pie chart and map of trash rating for Dorchester County streams sampled by the MBSS during 1995-97 and 2000-20048-154
8-90	Map of channelized sites, by type, for Dorchester County streams sampled by the MBSS during 2000-2004.....8-155
8-91	Map of sites with inadequate riparian buffers and buffer breaks for Dorchester County streams sampled by the MBSS during 2000-2004.....8-156
8-92	Pie charts and map of sites with eroded banks and instream bar formation for Dorchester County streams sampled by the MBSS during 2000-2004.....8-157
8-93	Pie chart and map of nitrate-nitrogen values for Dorchester County streams sampled by the MBSS during 1995-97 and 2000-20048-158
8-94	Pie chart and map of total phosphorus values for Dorchester County streams sampled by the MBSS during 2000-20048-159
8-95	Aquatic Heritage Biodiversity Ranking map for Dorchester County, by watershed8-160
FREDERICK COUNTY	
8-96	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Frederick County streams sampled by the MBSS during 1995-97 and 2000-20048-165
8-97	Physical Habitat Index pie chart and map of stream habitat quality for Frederick County streams sampled by the MBSS during 1995-97 and 2000-20048-168
8-98	Pie chart and map of trash rating for Frederick County streams sampled by the MBSS during 1995-97 and 2000-20048-169
8-99	Map of channelized sites, by type, for Frederick County streams sampled by the MBSS during 2000-2004.....8-170
8-100	Map of sites with inadequate riparian buffers and buffer breaks for Frederick County streams sampled by the MBSS during 2000-20048-171
8-101	Pie charts and map of sites with eroded banks and instream bar formation for Frederick County streams sampled by the MBSS during 2000-2004.....8-172
8-102	Pie chart and map of nitrate-nitrogen values for Frederick County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-173
8-103	Pie chart and map of total phosphorus values for Frederick County streams sampled by the MBSS during 2000-20048-174
8-104	Aquatic Heritage Biodiversity Ranking map for Frederick County, by watershed8-175
GARRETT COUNTY	
8-105	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Garrett County streams sampled by the MBSS during 1995-97 and 2000-20048-181
8-106	Physical Habitat Index pie chart and map of stream habitat quality for Garrett County streams sampled by the MBSS during 1995-97 and 2000-20048-184
8-107	Pie chart and map of trash rating for Garrett County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-185
8-108	Map of channelized sites, by type, for Garrett County streams sampled by the MBSS during 2000-2004.....8-186

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-109	Map of sites with inadequate riparian buffers and buffer breaks for Garrett County streams sampled by the MBSS during 2000-2004.....8-187
8-110	Pie charts and map of sites with eroded banks and instream bar formation for Garrett County streams sampled by the MBSS during 2000-2004.....8-188
8-111	Pie chart and map of nitrate-nitrogen values for Garrett County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-189
8-112	Pie chart and map of total phosphorus values for Garrett County streams sampled by the MBSS during 2000-2004.....8-190
8-113	Aquatic Heritage Biodiversity Ranking map for Garrett County, by watershed8-191
HARFORD COUNTY	
8-114	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Harford County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-197
8-115	Physical Habitat Index pie chart and map of stream habitat quality for Harford County streams sampled by the MBSS during 1995-97 and 2000-20048-200
8-116	Pie chart and map of trash rating for Harford County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-201
8-117	Map of channelized sites, by type, for Harford County streams sampled by the MBSS during 2000-2004.....8-202
8-118	Map of sites with inadequate riparian buffers and buffer breaks for Harford County streams sampled by the MBSS during 2000-2004.....8-203
8-119	Pie charts and map of sites with eroded banks and instream bar formation for Harford County streams sampled by the MBSS during 2000-2004.....8-204
8-120	Pie chart and map of nitrate-nitrogen values for Harford County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-205
8-121	Pie chart and map of total phosphorus values for Harford County streams sampled by the MBSS during 2000-2004.....8-206
8-122	Aquatic Heritage Biodiversity Ranking map for Harford County, by watershed8-207
HOWARD COUNTY	
8-123	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Howard County streams sampled by the MBSS during 1995-97 and 2000-20048-211
8-124	Physical Habitat Index pie chart and map of stream habitat quality for Howard County streams sampled by the MBSS during 1995-97 and 2000-20048-214
8-125	Pie chart and map of trash rating for Howard County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-215
8-126	Map of channelized sites, by type, for Howard County streams sampled by the MBSS during 2000-20048-216
8-127	Map of sites with inadequate riparian buffers and buffer breaks for Howard County streams sampled by the MBSS during 2000-2004.....8-217
8-128	Pie charts and map of sites with eroded banks and instream bar formation for Howard County streams sampled by the MBSS during 2000-2004.....8-218
8-129	Pie chart and map of nitrate-nitrogen values for Howard County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-219
8-130	Pie chart and map of total phosphorus values for Howard County streams sampled by the MBSS during 2000-2004.....8-220
8-131	Aquatic Heritage Biodiversity Ranking map for Howard County, by watershed.....8-221
KENT COUNTY	
8-132	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Kent County streams sampled by the MBSS during 1995-97 and 2000-20048-226
8-133	Physical Habitat Index pie chart and map of stream habitat quality for Kent County streams sampled by the MBSS during 1995-97 and 2000-20048-229
8-134	Pie chart and map of trash rating for Kent County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-230
8-135	Map of channelized sites, by type, for Kent County streams sampled by the MBSS during 2000-20048-231

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-136	Map of sites with inadequate riparian buffers and buffer breaks for Kent County streams sampled by the MBSS during 2000-2004.....8-232
8-137	Pie charts and map of sites with eroded banks and instream bar formation for Kent County streams sampled by the MBSS during 2000-2004.....8-233
8-138	Pie chart and map of nitrate-nitrogen values for Kent County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-234
8-139	Pie chart and map of total phosphorus values for Kent County streams sampled by the MBSS during 2000-2004.....8-235
8-140	Aquatic Heritage Biodiversity Ranking map for Kent County, by watershed.....8-236
MONTGOMERY COUNTY	
8-141	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Montgomery County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-241
8-142	Physical Habitat Index pie chart and map of stream habitat quality for Montgomery County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-244
8-143	Pie chart and map of trash rating for Montgomery County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-245
8-144	Map of channelized sites, by type, for Montgomery County streams sampled by the MBSS during 2000-2004.....8-246
8-145	Map of sites with inadequate riparian buffers and buffer breaks for Montgomery County streams sampled by the MBSS during 2000-2004.....8-247
8-146	Pie charts and map of sites with eroded banks and instream bar formation for Montgomery County streams sampled by the MBSS during 2000-2004.....8-248
8-147	Pie chart and map of nitrate-nitrogen values for Montgomery County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-249
8-148	Pie chart and map of total phosphorus values for Montgomery County streams sampled by the MBSS during 2000-2004.....8-250
8-149	Aquatic Heritage Biodiversity Ranking map for Montgomery County, by watershed.....8-251
PRINCE GEORGE'S COUNTY	
8-150	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Prince George's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-256
8-151	Physical Habitat Index pie chart and map of stream habitat quality for Prince George's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-259
8-152	Pie chart and map of trash rating for Prince George's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-260
8-153	Map of channelized sites, by type, for Prince George's County streams sampled by the MBSS during 2000-2004.....8-261
8-154	Map of sites with inadequate riparian buffers and buffer breaks for Prince George's County streams sampled by the MBSS during 2000-2004.....8-262
8-155	Pie charts and map of sites with eroded banks and instream bar formation for Prince George's County streams sampled by the MBSS during 2000-2004.....8-263
8-156	Pie chart and map of nitrate-nitrogen values for Prince George's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-264
8-157	Pie chart and map of total phosphorus values for Prince George's County streams sampled by the MBSS during 2000-2004.....8-265
8-158	Aquatic Heritage Biodiversity Ranking map for Prince George's County, by watershed.....8-266
QUEEN ANNE'S COUNTY	
8-159	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Queen Anne's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-270
8-160	Physical Habitat Index pie chart and map of stream habitat quality for Queen Anne's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-273
8-161	Pie chart and map of trash rating for Queen Anne's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-274

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-162	Map of channelized sites, by type, for Queen Anne's County streams sampled by the MBSS during 2000-2004.....8-275
8-163	Map of sites with inadequate riparian buffers and buffer breaks for Queen Anne's County streams sampled by the MBSS during 2000-2004.....8-276
8-164	Pie charts and map of sites with eroded banks and instream bar formation for Queen Anne's County streams sampled by the MBSS during 2000-2004.....8-277
8-165	Pie chart and map of nitrate-nitrogen values for Queen Anne's County streams sampled by the MBSS during 1995-97 and 2000-20048-278
8-166	Pie chart and map of total phosphorus values for Queen Anne's County streams sampled by the MBSS during 2000-20048-279
8-167	Aquatic Heritage Biodiversity Ranking map for Queen Anne's County, by watershed.....8-280
SOMERSET COUNTY	
8-168	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Somerset County streams sampled by the MBSS during 1995-97 and 2000-20048-284
8-169	Physical Habitat Index pie chart and map of stream habitat quality for Somerset County streams sampled by the MBSS during 1995-97 and 2000-20048-287
8-170	Pie chart and map of trash rating for Somerset County streams sampled by the MBSS during 1995-97 and 2000-20048-288
8-171	Map of channelized sites, by type, for Somerset County streams sampled by the MBSS during 2000-20048-289
8-172	Map of sites with inadequate riparian buffers and buffer breaks for Somerset County streams sampled by the MBSS during 2000-2004.....8-290
8-173	Pie charts and map of sites with eroded banks and instream bar formation for Somerset County streams sampled by the MBSS during 2000-2004.....8-291
8-174	Pie chart and map of nitrate-nitrogen values for Somerset County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-292
8-175	Pie chart and map of total phosphorus values for Somerset County streams sampled by the MBSS during 2000-2004.....8-293
8-176	Aquatic Heritage Biodiversity Ranking map for Somerset County, by watershed.....8-294
ST MARY'S COUNTY	
8-177	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for St. Mary's County streams sampled by the MBSS during 1995-97 and 2000-20048-299
8-178	Physical Habitat Index pie chart and map of stream habitat quality for St. Mary's County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-302
8-179	Pie chart and map of trash rating for St. Mary's County streams sampled by the MBSS during 1995-97 and 2000-20048-303
8-180	Map of channelized sites, by type, for St. Mary's County streams sampled by the MBSS during 2000-20048-304
8-181	Map of sites with inadequate riparian buffers and buffer breaks for St. Mary's County streams sampled by the MBSS during 2000-2004.....8-305
8-182	Pie charts and map of sites with eroded banks and instream bar formation for St. Mary's County streams sampled by the MBSS during 2000-2004.....8-306
8-183	Pie chart and map of nitrate-nitrogen values for St. Mary's County streams sampled by the MBSS during 1995-97 and 2000-20048-307
8-184	Pie chart and map of total phosphorus values for St. Mary's County streams sampled by the MBSS during 2000-20048-308
8-185	Aquatic Heritage Biodiversity Ranking map for St. Mary's County, by watershed.....8-309
TALBOT COUNTY	
8-186	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Talbot County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-314
8-187	Physical Habitat Index pie chart and map of stream habitat quality for Talbot County streams sampled by the MBSS during 1995-97 and 2000-20048-317
8-188	Pie chart and map of trash rating for Talbot County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-318

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-189	Map of channelized sites, by type, for Talbot County streams sampled by the MBSS during 2000-2004.....8-319
8-190	Map of sites with inadequate riparian buffers and buffer breaks for Talbot County streams sampled by the MBSS during 2000-2004.....8-320
8-191	Pie charts and map of sites with eroded banks and instream bar formation for Talbot County streams sampled by the MBSS during 2000-2004.....8-321
8-192	Pie chart and map of nitrate-nitrogen values for Talbot County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-322
8-193	Pie chart and map of total phosphorus values for Talbot County streams sampled by the MBSS during 2000-2004.....8-323
8-194	Aquatic Heritage Biodiversity Ranking map for Talbot County, by watershed8-324
WASHINGTON COUNTY	
8-195	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Washington County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-328
8-196	Physical Habitat Index pie chart and map of stream habitat quality for Washington County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-331
8-197	Pie chart and map of trash rating for Washington County streams sampled by the MBSS during 1995-97 and 2000-20048-332
8-198	Map of channelized sites, by type, for Washington County streams sampled by the MBSS during 2000-2004....8-333
8-199	Map of sites with inadequate riparian buffers and buffer breaks for Washington County streams sampled by the MBSS during 2000-2004.....8-334
8-200	Pie charts and map of sites with eroded banks and instream bar formation for Washington County streams sampled by the MBSS during 2000-2004.....8-335
8-201	Pie chart and map of nitrate-nitrogen values for Washington County streams sampled by the MBSS during 1995-97 and 2000-20048-336
8-202	Pie chart and map of total phosphorus values for Washington County streams sampled by the MBSS during 2000-20048-337
8-203	Aquatic Heritage Biodiversity Ranking map for Washington County, by watershed8-338
WICOMICO COUNTY	
8-204	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Wicomico County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-343
8-205	Physical Habitat Index pie chart and map of stream habitat quality for Wicomico County streams sampled by the MBSS during 1995-97 and 2000-20048-346
8-206	Pie chart and map of trash rating for Wicomico County streams sampled by the MBSS during 1995-97 and 2000-20048-347
8-207	Map of channelized sites, by type, for Wicomico County streams sampled by the MBSS during 2000-2004.....8-348
8-208	Map of sites with inadequate riparian buffers and buffer breaks for Wicomico County streams sampled by the MBSS during 2000-20048-349
8-209	Pie charts and map of sites with eroded banks and instream bar formation for Wicomico County streams sampled by the MBSS during 2000-2004.....8-350
8-210	Pie chart and map of nitrate-nitrogen values for Wicomico County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-351
8-211	Pie chart and map of total phosphorus values for Wicomico County streams sampled by the MBSS during 2000-2004.....8-352
8-212	Aquatic Heritage Biodiversity Ranking map for Wicomico County, by watershed8-353
WORCESTER COUNTY	
8-213	Benthic Index of Biotic Integrity and Fish Index of Biotic Integrity pie charts and map of stream health for Worcester County streams sampled by the MBSS during 1995-97 and 2000-2004.....8-359
8-214	Physical Habitat Index pie chart and map of stream habitat quality for Worcester County streams sampled by the MBSS during 1995-97 and 2000-20048-362
8-215	Pie chart and map of trash rating for Worcester County streams sampled by the MBSS during 1995-97 and 2000-20048-363
8-216	Map of channelized sites, by type, for Worcester County streams sampled by the MBSS during 2000-2004.....8-364

LIST OF FIGURES (CONTINUED)

Figure No.	Page
8-217 Map of sites with inadequate riparian buffers and buffer breaks for Worcester County streams sampled by the MBSS during 2000-2004	8-365
8-218 Pie charts and map of sites with eroded banks and instream bar formation for Worcester County streams sampled by the MBSS during 2000-2004.....	8-366
8-219 Pie chart and map of nitrate-nitrogen values for Worcester County streams sampled by the MBSS during 1995-97 and 2000-2004.....	8-367
8-220 Pie chart and map of total phosphorus values for Worcester County streams sampled by the MBSS during 2000-2004.....	8-368
8-221 Aquatic Heritage Biodiversity Ranking map for Worcester County, by watershed	8-369

8.1 INTRODUCTION

This report volume about stream conditions in Maryland's counties is part of a series of documents that details findings of the Maryland Biological Stream Survey (MBSS, or the Survey). It is particularly important because local-scale land planning decisions have a pronounced effect on stream and watershed conditions. Most volumes in the series use data from the 2000-2004 Survey. However, some documents, including this volume, utilize data from 1994 to 2004 and incorporate data from other sources as well.

The audience for this volume includes all those who have an interest in stream and watershed conditions in their county. The information contained here should help policymakers, planners, and others justify and guide natural resource conservation and restoration efforts in Maryland. In addition, the findings presented here may help guide other restoration efforts into the most needed areas, especially those associated with the Chesapeake Bay. In the interest of brevity, only select variables are presented in this volume, but county estimates can be calculated for all variables measured by the MBSS. In the interest of clarity, a **Glossary** of selected terms is provided at the back of this volume.

DATA FOR THIS REPORT:

A number of data sources were utilized for this volume, including data from the core MBSS program, county data that were submitted in response to our solicitation, and data from the volunteer component of MBSS, Stream Waders. The probability-based design of the core MBSS allowed for estimation of various aspects of condition with a known degree of confidence. County data augmented these findings, and Stream Waders data helped fill in gaps where no MBSS samples were collected. The next revision of this report will incorporate County data to a greater extent.

Every type of data collection has limitations and advantages, and the data used here are no exception. For all monitoring programs, sample size determines what can be concluded about a given geographic area, and the demand for information is often at a scale finer than can be accommodated by available data. Even with a minimum of ten sites in a watershed, confidence intervals about the estimates of condition are often quite high. One limitation more specific to MBSS data is that water chemistry samples are collected at a single point in time and in some cases may not capture the prevailing conditions at the site. In general, however, strong, consistent relationships between variables such as nitrate-nitrogen and percent agriculture in the upstream catchment confirm that the MBSS method of data collection does have utility. Another limitation of the MBSS is that subtle trends in condition in small geographic areas are difficult to detect because only Sentinel sites are revisited each year. However, if improvements in condition occur across larger areas, these trends will likely be detected by the MBSS.

One limitation of Stream Waders data is the uncertainty about whether samples were collected properly by volunteers, whether the correct coordinates and site information was recorded, etc. However, Quality Control visits and duplicate sampling by professionals have repeatedly demonstrated that Stream Waders data is of generally high quality. An additional limitation is that non-randomly selected sites are not generally useful for condition estimates. A major advantage of Stream Waders data is high spatial density. Because of this density, at least general statements can be made about the condition of many small watersheds that were never sampled before.

Finally, there are limitations with direct incorporation of data collected by county agencies, because no such agency has a sampling design identical to the core MBSS. The difficulties of incorporation are formidable and costly, but do enhance the ability of counties to draw conclusions from the data.

WHO CAN USE THIS DOCUMENT:

Citizens and Local Environmental Groups - to better understand and provide input on conservation and restoration initiatives

Landowners - to make informed decisions about long-term conservation and restoration on their property

Local Governments - to incorporate sound, targeted conservation practices, policies, and zoning

Educational Institutions - to provide a conservation education tool for use by students in each county

This volume contains a chapter summarizing conditions in all counties, followed by a chapter for each individual county and one for Baltimore City. Other volumes in this series also contain information that may be of use to the reader, especially the volumes on biodiversity (Volume 12) and riparian buffers (Volume 10). To limit the size and complexity of this volume and increase readability, all methods used to prepare and analyze data for this volume are presented in: 2000-2004 Maryland Biological Stream Survey Laboratory, Field, and Analytical Methods: Volume 6. This volume and others can be downloaded from <http://www.dnr.Maryland.gov/streams/pubs/>.

8.2 COUNTY COMPARISONS

8.2.1 Biotic Integrity

The Combined Biotic Index (CBI) was developed to provide a single measure of stream health for Maryland streams. The CBI is derived from the Fish Index of Biotic Integrity (FIBI) and the benthic macroinvertebrate Index of Biotic Integrity (BIBI). Among all counties in Maryland during 2000-2004, Queen Anne's County had the highest Combined Biotic Index (CBI) score (3.49), with an overall rating of Fair (Table 8-1). Other counties in the top five included Charles, Carroll, St. Mary's, and Talbot. In contrast, Baltimore City had the lowest average CBI (2.18), with an overall stream integrity rating of Poor. The next lowest counties were Somerset, Dorchester, Anne Arundel, and Washington.

For the Fish Index of Biotic Integrity (FIBI), Carroll County had the highest mean score (3.62), with an overall rating of Fair (Table 8-1). The next highest four counties were Howard, Queen Anne's, Cecil, and Kent County. Caroline County had the lowest mean FIBI score (2.23), followed by Calvert, Baltimore City, Dorchester, and Anne Arundel. Overall, 13 counties and Baltimore City had an average FIBI rating of Poor, ten counties rated Fair, and no county had a mean FIBI score in the Good range.

The overall results for the Benthic Index of Biotic Integrity (BIBI) were generally similar to the FIBI. Fourteen counties rated in the fair category, ten counties and Baltimore City were rated in the Poor category (Table 8-1). Among individual counties, St. Mary's County had the highest mean BIBI score, and Baltimore City had the lowest mean score. Interestingly, the highest four counties for mean BIBI (Charles, Talbot, St. Mary's, and Queen Anne's) are all located in the Coastal Plain region.

In general, urban or urbanizing counties had higher scores for fish than for benthic macroinvertebrates. A possible explanation for this phenomenon is that fish are more mobile than benthos, and thus better able to move to refugia during adverse conditions such as high flows during summer thunderstorms. In contrast, mean scores for fish in Calvert and Anne Arundel Counties were considerably lower than mean benthic scores. The networks of many streams in these counties are relatively small and isolated from recolonization by fish because of the salinity in Chesapeake Bay. Thus, historical problems of a watershed-wide nature (e.g., clearcutting) could result in elimination of some fish species because there was no chance for recolonization. In contrast, at least some benthic macroinvertebrates are capable of movement to neighboring watersheds (via flying as adults).

The BIBI and FIBI results presented in this report have been validated as being able to consistently distinguish between degraded and reference conditions (Southerland et al., 2005). However, like any tool, indices of biotic integrity may not perform at a uniform level throughout the range of conditions where they are used. In addition, different taxonomic groups may not respond in the same way to the myriad number of stressors to Maryland streams. Thus, some differences in ratings between assemblages should be expected.

8.2.2 Chemical Conditions

As described earlier, MBSS water chemistry data represent grab samples from a single point in time, and thus may not capture prevailing conditions at a site or

IBI SCORES AND RATING TRANSLATION

The Index of Biotic Integrity, or IBI, was developed to help rate the health of streams, including their water quality. The IBIs developed for Maryland (see Southerland *et al.*, 2005) are based on a series of reference sites that represent some of the best remaining stream habitats in the state. However, because most areas of Maryland have been subject to extensive human disturbance and there are no streams left that even approach pristine conditions, IBIs tend to overrate the actual quality of streams. To make it easier to interpret, numeric ratings are grouped into descriptive categories. In the MBSS, ratings are Good, Fair, Poor, and Very Poor. The lack of an Excellent rating is a recognition of the fact that the reference conditions used to develop the IBIs often have significant human perturbation associated with them.

seasonal/event driven changes. However, these data have been useful in describing overall patterns as well as illustrating specific water quality problems. For example, it is highly likely that if low dissolved oxygen levels are observed at a site on one occasion, low levels have been occurring at other times as well. Further, even short duration exposure to some conditions can profoundly alter the biological community.

8.2.2.1 Dissolved Oxygen

During the 2000-2004 MBSS, several Eastern Shore counties were observed to have a relatively high proportion of stream miles with dissolved oxygen (DO) problems. Although the Maryland regulatory criterion for low DO is 5 mg/l, 3 mg/l was used as the threshold for this report (a DO of 5 mg/l may be within the range of natural variability). Counties with low DO included: Somerset (38% of stream miles), Dorchester (26%), Caroline (26%), and Baltimore City (23%) (Table 8-2).

Table 8-1. Mean biological integrity scores for 4 th order and smaller Maryland streams, by county, based on results from the 2000-2004 MBSS									
COUNTY	Combined Biotic Index (CBI)			Fish Index of Biotic Integrity (FIBI)			Benthic Index of Biotic Integrity (BIBI)		
	Mean	SE	Rank	Mean	SE	Rank	Mean	SE	Rank
Allegany	3.05	0.10	11	2.63	0.15	19	3.48	0.11	5
Anne Arundel	2.66	0.13	21	2.48	0.22	20	2.81	0.14	16
Baltimore	2.92	0.13	13	2.35	0.37	22	3.05	0.15	13
Baltimore City	2.18	0.28	24	2.76	0.17	18	2.12	0.07	24
Calvert	2.82	0.21	15	2.31	0.30	23	3.34	0.21	6
Caroline	2.80	0.17	16	2.23	0.20	24	3.22	0.17	11
Carroll	3.42	0.11	3	3.62	0.15	1	3.26	0.12	8
Cecil	3.33	0.18	7	3.43	0.20	4	3.25	0.21	9
Charles	3.43	0.11	2	3.03	0.15	10	3.83	0.12	2
Dorchester	2.54	0.19	22	2.42	0.34	21	2.61	0.22	21
Frederick	2.68	0.12	19	2.86	0.19	12	2.52	0.09	23
Garrett	3.08	0.11	10	2.85	0.12	14	3.33	0.12	7
Harford	3.19	0.08	8	3.28	0.13	8	3.18	0.09	12
Howard	3.34	0.09	6	3.45	0.13	2	3.23	0.14	10
Kent	3.17	0.15	9	3.41	0.21	5	3.00	0.16	14
Montgomery	2.89	0.10	14	3.16	0.16	9	2.65	0.09	19
Prince Georges	2.78	0.11	17	2.83	0.15	15	2.72	0.11	18
Queen Annes	3.49	0.20	1	3.44	0.24	3	3.50	0.21	4
Somerset	2.50	0.17	23	3.30	0.29	6	2.54	0.19	22
St Marys	3.40	0.11	4	2.86	0.15	13	3.89	0.13	1
Talbot	3.35	0.34	5	2.94	0.45	11	3.75	0.24	3
Washington	2.67	0.12	20	2.78	0.17	17	2.62	0.11	20
Wicomico	2.99	0.17	12	3.30	0.17	7	2.93	0.19	15
Worcester	2.72	0.19	18	2.82	0.29	16	2.78	0.21	17

Nine counties had no stream miles with very low DO conditions.

In the Coastal Plain region of Maryland, streams with high levels of dissolved organic carbon (DOC) tended to have low DO levels (Figure 8-1). Although the naturally poor light penetration in this tea-stained water is a likely contributing factor to the lower DO, many of the watersheds containing high DOC also have nutrient loads that are some of the highest in the state.

8.2.2.2 Nitrogen

Among all counties and Baltimore City, streams in Caroline County had the highest mean value for Total Nitrogen (6.24 mg/l), and Charles County streams had the lowest mean Total Nitrogen value (0.74 mg/l; Table 8-2). Caroline County also had the highest mean nitrate-nitrogen levels (5.51 mg/l), followed by Dorchester County (5.20 mg/l). The lowest nitrate-nitrogen level was observed in Somerset County streams (0.40 mg/l). There

was a strong relationship between mean nitrate-nitrogen levels for each county and the percentage of that county harvested as cropland (Figure 8-2).

For ammonia, the county with the highest mean value was Kent County (0.22 mg/l), and the lowest levels were observed in Allegany and Garrett counties (0.01 mg/l).

8.2.2.3 Phosphorous

Anne Arundel (0.010 mg/l) and Garrett (0.013 mg/l) county streams had the lowest mean levels of Total Phosphorous observed during 2000-2004 (Table 8-2). In contrast, Worcester (0.125 mg/l) and Kent Counties (0.127 mg/l) had the highest mean levels in the state during 2000-2004. The total number of chickens reported for each county was significantly related to the mean Total Phosphorus concentrations in that county (Figure 8-3).

Table 8-2. Mean values for key chemical constituents of 4th order and smaller Maryland streams, by county, based on results from the 2000-2004 MBSS

	Total Nitrogen			Nitrate-N			Ammonia			Total Phosphorus			Dissolved Oxygen			pH			Acid Neutralizing Capacity (ANC)		
COUNTY	Mean (mg/l)	SE	Rank	Mean (mg/l)	SE	Rank	Mean (mg/l)	SE	Rank	Mean (mg/l)	SE	Rank	% Stream miles < 3 mg/l	SE	Rank	Mean	SE	Rank	Mean (mg/l)	SE	Rank
Allegany	0.75	0.06	2	0.65	0.06	6	0.01	0.00	1	0.010	0.001	1	0.0	0.0	1	6.59	0.13	17	440.15	50.17	13
Anne Arundel	1.08	0.09	8	0.81	0.08	7	0.06	0.01	17	0.049	0.007	14	5.9	2.9	6	7.15	0.05	10	427.26	70.79	15
Baltimore	2.43	0.16	12	2.29	0.16	12	0.03	0.01	8	0.042	0.015	10	0.0	0.0	1	7.65	0.05	2	1368.95	191.54	3
Baltimore City	1.98	0.37	10	1.76	0.35	10	0.03	0.00	6	0.040	0.016	8	23.4	16.8	13	7.66	0.03	1	1393.13	70.64	2
Calvert	0.76	0.14	4	0.61	0.14	5	0.06	0.01	15	0.077	0.012	22	0.0	0.0	1	6.98	0.10	11	558.73	94.79	9
Caroline	6.24	0.43	24	5.51	0.43	24	0.06	0.02	18	0.049	0.006	13	25.6	9.2	14	7.35	0.15	7	550.76	63.29	10
Carroll	3.78	0.32	20	3.42	0.28	20	0.14	0.10	23	0.046	0.019	11	0.0	0.0	1	6.31	0.08	19	163.96	18.92	22
Cecil	2.75	0.25	16	2.40	0.25	14	0.05	0.01	14	0.062	0.011	19	4.6	4.6	4	6.23	0.11	21	195.00	22.34	20
Charles	0.74	0.08	1	0.48	0.06	2	0.07	0.02	20	0.040	0.006	9	7.3	3.1	7	7.28	0.05	8	608.94	55.20	7
Dorchester	5.20	0.68	23	4.59	0.66	23	0.05	0.01	11	0.075	0.019	21	26.1	8.0	15	6.08	0.12	22	198.26	39.09	19
Frederick	2.69	0.26	13	2.43	0.24	15	0.05	0.01	13	0.053	0.010	15	0.0	0.0	1	7.44	0.09	5	905.96	109.14	4
Garrett	0.96	0.08	6	0.86	0.08	8	0.01	0.00	2	0.013	0.001	2	0.0	0.0	1	6.64	0.10	16	177.66	28.08	21
Harford	2.90	0.19	17	2.57	0.18	18	0.02	0.00	5	0.030	0.003	4	3.8	2.1	3	7.16	0.05	9	423.51	26.40	16
Howard	2.72	0.17	14	2.46	0.16	16	0.02	0.00	4	0.037	0.019	6	0.0	0.0	1	7.40	0.07	6	743.83	65.85	6
Kent	3.77	0.58	19	2.97	0.53	19	0.22	0.11	24	0.117	0.020	23	8.2	5.3	9	6.70	0.08	13	465.60	59.77	11
Montgomery	2.30	0.19	11	2.10	0.19	11	0.03	0.01	7	0.027	0.003	3	0.0	0.0	1	7.47	0.08	4	743.93	65.66	5
Prince Georges	0.92	0.07	5	0.61	0.05	4	0.09	0.03	21	0.048	0.005	12	4.9	3.2	5	6.88	0.07	12	592.50	80.51	8
Queen Annes	4.26	0.38	22	3.68	0.36	22	0.06	0.03	19	0.056	0.006	17	3.2	2.8	2	6.65	0.09	15	429.26	41.76	14
Somerset	1.01	0.14	7	0.40	0.12	1	0.05	0.01	12	0.060	0.028	18	38.3	8.5	16	6.53	0.11	18	338.83	80.09	17
St Marys	0.76	0.09	3	0.57	0.09	3	0.04	0.01	9	0.038	0.007	7	12.2	5.7	10	5.21	0.11	24	49.44	12.09	24
Talbot	4.15	0.71	21	3.48	0.71	21	0.04	0.01	10	0.055	0.009	16	22.5	15.9	12	6.67	0.11	14	457.14	48.58	12
Washington	2.74	0.33	15	2.49	0.28	17	0.02	0.01	3	0.037	0.006	5	0.0	0.0	1	7.59	0.10	3	1985.02	241.79	1
Wicomico	3.07	0.29	18	2.40	0.29	13	0.06	0.01	16	0.067	0.017	20	20.3	8.6	11	6.26	0.13	20	222.69	30.33	18
Worcester	1.91	0.41	9	1.35	0.37	9	0.11	0.08	22	0.125	0.049	24	7.6	7.5	8	5.84	0.21	23	163.21	35.16	23




Figure 8-1. Dissolved organic carbon (DOC) levels vs. dissolved oxygen (DO) from the 2000-2004 MBSS




Figure 8-2. Mean nitrate-nitrogen levels from the 2000-2004 MBSS, by county, and the percentage of that county harvested as cropland (crop data from USDA 2002)




Figure 8-3. Mean Total Phosphorous levels from the 2000-2004 MBSS, by county, and the number of chickens raised in the county (poultry data from USDA 2002)

NATURAL NUTRIENT LEVELS IN MARYLAND STREAMS

A commonly asked question when nutrients in streams are discussed is: what are background or 'natural' levels of nutrient concentration in streams? As essentially all of Maryland has been logged, affected by atmospheric deposition of nitrogen, and otherwise disturbed at one time or another, this question is difficult to answer. However, to gain some insight into this question, all sites in the 1995-2004 dataset with > 90% forest in the upstream catchment were analyzed separately. The mean, median, 90th percentile, and 10th percentile for nitrogen and phosphorus species are shown below, along with chloride, a general indicator of human disturbance. It should be noted that because of the level of human disturbance in many soil types and geologic strata, some areas of Maryland may not be accurately characterized by the information presented. However, as there is a strong trend of decreasing nutrient levels with increases in percent forest in the corresponding watershed, it follows that nutrient levels in old growth, undisturbed forests throughout Maryland would be lower than the best sites that remain today.

Summary of nutrient parameters for MBSS water quality collected during spring baseflow in first through third order streams for all Maryland sites (both rounds) with greater than a 90% forested watershed (units = mg/L).

Parameter	Mean	Median	Lower Quartile	Upper Quartile	10% Percentile	90% Percentile	N
Nitrate-N	0.47	0.31	0.14	0.61	0.06	1.14	187
Nitrite-N	0.0011	0.0004	0.0004	0.001	0.0004	0.0025	107
Ammonia-N	0.012	0.0044	0.0024	0.015	0.002	0.027	114
Total N	0.59	0.46	0.22	0.73	0.12	1.45	134
Total P	0.012	0.008	0.006	0.015	0.004	0.021	134
Ortho-PO ₄	0.003	0.0007	0.0007	0.0035	0.0006	0.0078	126
Chloride	5.98	3.04	1.23	6.28	1.01	10.5	134

8.2.2.4 pH and Acid Neutralizing Capacity (ANC)

The streams with the highest mean pH values (uncorrected for log scale) in the state were in Baltimore City (7.66) and Baltimore County (7.65) (Table 8-2). In contrast, the lowest mean values were observed in St. Mary's (5.2) and Worcester (5.84) streams. In terms of buffering capacity against acid inputs, Washington County had the highest mean ANC value (1985 µeq/l), and Somerset County had the lowest mean ANC (49 µeq/l). The generally elevated pH and buffering capacity observed in Baltimore City and County are most likely due to the effects of urbanization, and the low pH and buffering capacity on the lower eastern shore may be due to naturally acidic conditions that have been exacerbated by the continuing effects of acid deposition and the application of nitrogen fertilizer.

Streams with ANC less than 0 µeq/l are acidic and very poorly buffered. Streams with ANC between 0 and 200 µeq/l are only moderately buffered and may periodically have low pH levels during rain or snowmelt events. Those streams with ANC greater than 200 µeq/l are well-buffered.

8.2.3 Physical Conditions

8.2.3.1 Overall Conditions

Physical conditions play a key role in determining stream health. To provide an overall indication of stream habitat quality, the MBSS developed a multi-metric Physical Habitat Index (PHI). As with the Indices of Biotic Integrity for fish and benthic macroinvertebrates, the PHI was developed using a set of defined reference and degraded conditions and an independent set of data was used for validation purposes. Specific components of the index vary by region, but the core metrics include remoteness, epifaunal substrate, bank stability, and shading.

Of all Maryland counties and Baltimore City, Allegany and Garrett counties had the highest mean PHI values (77 on a 100 point scale) during 2000-2004 (Table 8-3). In contrast, Baltimore City (50/100) and Cecil County (63/100) had the lowest mean habitat values in the state during that timeframe.

COUNTY	Physical Habitat Index (PHI)			Channelized ¹			Trash Rating			Bank Erosion			Riparian Buffer			Riparian Buffer Breaks		
	Mean Score (0-100)	SE	Rank	% Stream miles	SE	Rank	Mean Score (0-20)	SE	Rank	% Stream miles Optimal ²	SE	Rank	% Stream miles no buffer	SE	Rank	% Stream miles with Severe Buffer Breaks ³	SE	Break Rank
Allegany	76.52	1.77	2	17.2	4.3	11	16.3	0.5	14	88.0	3.8	3	6.6	2.8	14	3.8	2.0	3
Anne Arundel	68.39	1.91	16	19.9	4.8	13	11.9	0.7	3	30.5	6.9	22	2.7	1.7	10	12.4	4.8	17
Baltimore	64.34	1.90	22	21.8	5.5	15	12.9	0.6	4	25.5	5.9	23	7.8	3.9	17	14.6	4.8	21
Baltimore City	50.21	2.31	24	40.5	15.0	18	5.2	1.8	1	81.2	13.2	6	9.3	9.2	19	43.8	14.6	24
Calvert	66.85	3.74	21	9.0	9.0	3	16.4	1.0	16	41.0	13.9	16	0.0	0.0	1	9.0	9.0	13
Caroline	70.90	1.55	8	58.3	9.0	22	16.6	0.6	19	73.0	10.4	9	2.1	2.0	8	4.2	4.1	4
Carroll	69.38	1.54	12	13.1	5.4	6	16.2	0.3	13	34.7	9.2	20	10.0	4.1	20	10.5	4.1	15
Cecil	62.82	2.34	23	14.2	6.9	8	14.3	0.6	7	42.2	8.6	15	11.2	6.2	22	5.4	3.6	6
Charles	75.67	1.18	3	10.5	3.8	4	16.3	0.4	15	57.1	5.5	12	0.0	0.0	2	0.9	0.9	2
Dorchester	69.06	2.18	14	57.1	10.1	21	16.5	0.8	17	82.4	8.2	5	0.0	0.0	3	10.3	5.4	14
Frederick	70.37	2.00	11	14.6	5.2	9	15.6	0.5	10	62.7	7.2	11	11.1	4.2	21	12.8	5.0	18
Garrett	77.02	1.86	1	6.6	2.8	2	17.6	0.3	24	88.9	4.2	1	7.7	3.4	15	7.4	3.3	12
Harford	68.26	1.33	17	18.8	3.9	12	13.0	0.4	5	55.5	5.5	13	5.9	2.5	13	6.7	2.8	10
Howard	71.52	1.79	6	16.5	5.5	10	14.4	0.5	8	19.5	5.5	24	4.8	3.4	11	6.8	4.0	11
Kent	67.80	2.14	18	11.2	8.0	5	16.9	0.5	21	38.1	8.2	17	1.2	1.2	7	6.6	6.6	9
Montgomery	70.75	1.17	9	21.6	5.2	14	13.8	0.5	6	35.3	6.6	19	7.7	3.6	16	10.5	4.1	16
Prince Georges	67.41	1.88	20	31.4	5.6	16	10.9	0.6	2	51.9	7.1	14	2.1	2.1	9	13.5	4.0	19
Queen Annes	67.45	2.66	19	43.1	10.7	19	17.1	0.4	23	64.5	11.4	10	0.0	0.0	4	4.5	4.5	5
Somerset	70.57	1.86	10	95.5	4.5	24	16.6	0.5	18	84.3	7.9	4	5.4	5.4	12	5.4	5.4	7
St Marys	72.41	1.82	5	3.9	2.9	1	16.2	0.6	12	36.3	9.2	18	1.0	1.0	6	5.5	3.7	8
Talbot	71.29	1.84	7	33.6	19.3	17	16.0	0.5	11	33.7	11.3	21	0.0	0.0	5	0.0	0.0	1
Washington	69.33	2.51	13	13.2	5.1	7	16.6	0.4	20	77.0	6.7	8	20.1	6.0	24	13.5	4.3	20
Wicomico	72.67	2.21	4	52.1	10.0	20	14.6	0.8	9	88.0	6.6	2	19.2	7.0	23	19.2	7.0	23
Worcester	68.91	1.73	15	67.9	9.4	23	17.0	0.6	22	77.4	9.6	7	9.0	5.0	18	14.8	5.4	22
¹ Estimate based only on sites that were completely channelized ² Optimal bank erosion defined as bank erosion ≥ 16 on a 16-20 scale. ³ Estimate based on 75m sites that had discontinuity in the riparian buffer that was judged to have a significant negative impact on the stream during storm events or baseflow conditions (e.g., raw sewage entry)																		

8.2.3.2 Trash

Human refuse in and along streams is unsightly but may have little influence on chemical and physical conditions. However, trash along streams is an indication that other, more toxic contaminants such as oil and antifreeze have been dumped. As would be expected, urban counties had lower trash rating scores than agricultural counties (Table 8-3). Baltimore City received the lowest mean rating (5 on a 20 point scale with lower scores indicating more trash), followed by Prince George's, Anne Arundel, and Baltimore counties. The least amount of human refuse on average along or in streams was found in Garrett (18/20) and Somerset counties (17/20).

8.2.3.3 Channelization

In Maryland, stream channels are moved and/or hardened for a number of reasons, including rapid routing of flood waters, protection of property and transportation infrastructure, and lowering of water tables to facilitate the use of land for agriculture. Unfortunately, channelized streams have reduced habitat quality and are less able to retain suspended sediments and retain and process nutrients. The ultimate result is movement of sediment and excess nutrients into estuaries such as Chesapeake Bay, where they cause 'dead zones' that are devoid or

depleted of dissolved oxygen and degraded habitats for submerged aquatic vegetation. Dead zones, in turn cause significant damage to commercial and recreational fisheries. During the 2000-2004 MBSS, the estimated percent of non-tidal stream miles that were channelized varied from 4% in St. Mary's County to 96% in Somerset County (Table 8-3). Channelization was also a dominant stream feature in two other Eastern Shore counties: Worcester (68%) and Dorchester (57%). Among the counties with substantial urban land use, channelization varied from 20% in Anne Arundel County to 41% in Baltimore City.

Among various types of channelization, only ditches were found in more than 50% of the stream miles in a county (Table 8-4). This occurred in Somerset (96%), Worcester (65%), Caroline (58%), and Dorchester (54%) counties. Only four counties and Baltimore City had no ditched streams. In contrast, concrete channels were most extensive in more urban jurisdictions. Baltimore City had an estimated 28% of stream miles in concrete, followed by Baltimore County (11%), Prince George's (9%), and Montgomery (6%). Concrete channels provide exceptionally poor habitat for stream organisms, do not dissipate energy, and route water rapidly downstream, exacerbating flooding.

County	Concrete	Ditched	Gabion	Pipe	RipRap	None
Allegany	4.6	1.3	1.1	3.4	4.3	85.3
Anne Arundel	2.5	4.8	1.2	0.0	7.8	83.8
Baltimore	11.1	0.7	2.1	0.6	7.3	78.2
Baltimore City	28.0	0.0	6.3	0.0	6.3	59.5
Calvert	0.0	9.0	0.0	0.0	0.0	91.0
Caroline	0.0	58.3	0.0	0.0	0.0	41.7
Carroll	0.6	8.1	0.0	3.3	1.1	86.9
Cecil	2.6	0.0	0.0	2.6	9.0	85.8
Charles	0.0	5.0	0.0	4.8	0.6	89.5
Dorchester	0.0	53.8	0.0	0.0	0.0	46.2
Frederick	0.0	0.0	2.9	5.8	5.9	85.4
Garrett	1.4	0.0	1.2	1.5	2.5	93.4
Harford	3.2	7.7	0.0	2.6	5.3	81.2
Howard	1.6	2.6	0.0	4.5	7.8	83.5
Kent	6.4	4.7	0.0	0.0	0.0	88.8
Montgomery	5.8	0.0	0.0	2.0	13.9	78.3
Prince Georges	8.9	10.4	1.1	1.8	9.3	68.6
Queen Annes	4.5	36.0	0.0	2.6	0.0	56.9
Somerset	0.0	95.5	0.0	0.0	0.0	4.5
St Marys	0.0	3.9	0.0	0.0	0.0	96.1
Talbot	0.0	22.5	0.0	0.0	0.0	77.5
Washington	4.9	0.8	0.0	4.3	3.2	86.8
Wicomico	0.0	42.5	3.8	5.9	0.0	47.9
Worcester	0.0	65.0	0.0	0.0	0.0	35.0

Gabions, or rock-filled baskets, were much less extensive than either concrete channels or ditches. Baltimore City had the highest percentage of stream miles with gabions (6%), and only 7 of 23 Maryland counties and Baltimore City had gabions in any of their randomly selected MBSS sites. Rip-rap, in contrast, was most common in Montgomery County (14% of stream miles) and was documented in 14 of the 24 jurisdictions. Piped culverts were most common at MBSS sites in Wicomico and Frederick counties (6% of stream miles in each), but culverts occur throughout the state and are the most commonly used method of passing streams under roads.

8.2.3.4 Bank Erosion

Excessively eroded banks are a symptom of watershed-scale problems such as impervious surface-related runoff, unwise harvest of riparian vegetation, and stream elevation changes from improper culvert installation. The 2000-2004 MBSS data indicate that Howard County had the lowest percentage of stream miles with little to no evidence of bank erosion (20%). Other counties with relatively few intact stream banks included: Baltimore City (26%), Baltimore (31%), and Talbot (34%; Table 8-3). In contrast, nearly 90% of stream banks were rated as optimal for minimal bank erosion in Garrett, Allegany, and Wicomico counties.

8.2.4 Riparian Conditions

Riparian buffers play a key role in providing instream habitat, temperature control, organic matter inputs, and energy dispersal during storm events. In addition, they also filter excess nutrients and other contaminants before they reach the stream. Among all counties in Maryland, Washington County had the highest estimate of stream miles with no riparian buffers (20%), followed by Wicomico (19%), Cecil (11%) and Frederick (11%; Table 8-3). In contrast, no bufferless stream sites were observed in Calvert, Charles, Queen Anne's, Dorchester, and Talbot counties.

In cases where breaks in riparian buffers exist, the beneficial aspects of buffers can be greatly reduced or virtually eliminated. For the MBSS, breaks in the riparian buffer zone were defined as any short-circuiting of the existing vegetation that allowed water, sediment, or potential contaminants to flow directly into the stream without passing through or over a buffered area. Breaks were rated as minor or severe, and types included storm and tile drains, impervious surfaces and erosion gullies, crops, orchards and pastures, and roads and railroads. Of the counties with no totally unbuffered sites, all counties but Talbot had sites with severe breaks in existing buffers (Table 8-3). Other counties with less than 5% of stream miles with severe buffer breaks included: Allegany, Caroline, Charles, Kent, and Queen Anne's. Counties

with the highest occurrence of buffer breaks were Baltimore City (44%), Worcester (20%), Wicomico (15%), and Baltimore counties (15%). From these analyses, it is clear that buffer breaks are a potentially important limitation to the effectiveness of riparian buffers in protecting Maryland streams.

8.2.5 County Stressors

Identifying stressors is critical to the development of management actions by counties to restore and protect streams. In particular, counties have a key role in implementing Total Maximum Daily Loads (TMDLs) developed by the State to address streams impaired under Section 303(d) of the Clean Water Act. Counties are also key partners in active restoration programs for Maryland's streams, such as Maryland's Watershed Restoration Action Strategies (WRASs) and Chesapeake Bay Program Tributary Strategies. Although TMDL development and other activities rely on the identification of specific causes of degradation to guide actions, unless relevant information is available to reliably identify these stressors, funding, effort and political capital could be spent on problems that are not really prevalent at the watershed scale.

Merely identifying stressors, however, is still not sufficient to guide effective management actions for Maryland streams. Counties and other natural resource stewards must assess the relative degree of risk posed by different stressors at site, watershed, and regional scales. Only by comparing these risks and determining the cumulative impacts that are likely to result, can an effective stream restoration and protection strategy be implemented. To fully assess the threat from an individual stressor, the importance (or severity) and the prevalence (or extent) of the stressor must be known. The severity of each stressor was assessed based on the response of fish and benthic macroinvertebrate Indices of Biotic Integrity (IBI) scores to established thresholds.

Streams and their inhabitants are subject to myriad stressors, some of which may be nearly impossible to measure, co-occur with other stressors, or otherwise difficult to assess. To contribute to what is known about stressors and biological responses in Maryland, a total of ten stressors were identified from MBSS data and landscape-level data (see sidebar). For each stressor, the proportion of MBSS sites having poor fish or benthic IBI scores when the threshold for degradation was exceeded was divided by the proportion of sites with poor IBI scores, given stressor scores below the threshold for degradation (Figure 8-4). Applying this technique to stressors measured as part of the MBSS, the stressor that caused the greatest difference in IBI scores above and below the threshold was Acid Mine Drainage. By incorporating both severity and extent into stressor analysis, important insight can be gained as to which

Identifying stressors is critical to meeting Clean Water Act mandates and developing management actions that can restore or protect the desired condition of streams. Stressor identification, or the diagnosis of stream problems, is an emerging field that draws on the approaches of traditional risk assessment while using new metrics derived from more sophisticated monitoring data. Therefore, the MBSS is conducting analyses in this and other volumes to investigate which stressors are responsible for degradation of Maryland streams.

Stressors can be organized according to the five major determinants of biological integrity in aquatic ecosystems: water chemistry, energy source, habitat structure, flow regime, and biotic interactions. Water chemistry comprises acidity, dissolved oxygen, and contaminants. Energy source describes the size, abundance, and nutritional quality of food from both primary production and allochthonous inputs. Habitat structure encompasses physical features such as water depth, current velocity, substrate composition, and morphology of the stream channel. Flow regime refers to seasonal, annual, and altered patterns in water quantity and delivery. Biotic interactions include competition, predation, and parasitism, from both native and introduced species. The MBSS directly measures many of these stressors and ancillary information, such as land use, can be used to evaluate others. Some stressors, such as pesticides, currently are not considered in MBSS analyses. This volume includes analysis of ten stressors affecting Maryland streams: invasive fish and mussels, invasive plants, bank stability, acid mine drainage, acidic deposition, dissolved oxygen, high nitrate-nitrogen, channelization, no riparian buffer, and urban land use (see below). These 10 stressors are meant to be a representative but incomplete list. The thresholds of concern for each stressor were selected based on expert consensus and analyses to date on the MBSS data. In particular, stressor values that result in demonstratively lower fish or benthic IBI scores have been used as thresholds. Additional stressor analyses are being conducted with the MBSS data and thresholds may be revised in the future.

LIST OF STRESSORS AND THRESHOLDS USED IN THE 2000-2004 MBSS:

Urban	> 5%	Dissolved Oxygen	< 3 mg/L
Bank Stabling	Poor or Very Poor	Acid Mine Drainage	Present
Channelization	Present	Acid Deposition	Present
Riparian Buffer	0 m	Invasive Plants	Present
Nitrate-Nitrogen	> 5mg/L	Non-Native Aquatic Species (Fish or Bivalves)	Present

stressors have the strongest effect on streams statewide or within a county or watershed.

Because the MBSS sampling design is probability-based, the extent of any stressor that is measured can be estimated with a known degree of confidence. To determine the extent of each stressor in each county and Baltimore City, we estimated the percentage of stream miles with stressor scores above the threshold of degradation. Based on these estimates, the single most prevalent stressor affecting Maryland streams was the presence of non-native invasive plants (Figure 8-5). This stressor was present in 100% of stream miles for Baltimore, Cecil, Carroll, Howard, and Montgomery counties, and high in many other counties as well. Of the other stressors characterized by the MBSS, urban influence was greatest in Baltimore City and Anne Arundel County, while unstable banks were most extensive in Howard and Talbot counties. Channelization was most extensive in Somerset and Worcester county streams, and Washington County had the most stream

miles with no riparian buffer. Deleterious nitrate-nitrogen levels were most frequent in Caroline and Dorchester county streams, while low dissolved oxygen was most extensive in Somerset County. Acid Mine Drainage existed only in Allegany and Garrett Counties, while acid deposition effects were much more widespread and most extensive in St. Mary's and Garrett counties. And finally, non-native aquatic animals, including fish, occurred extensively in all Maryland counties but were most prevalent in Howard and Kent counties.

The extent of each stressor, combined with the severity estimate, provides useful information about the relative risk each stressor poses to streams within areas of interest such as counties. A more complete discussion of stressors and their relative risks to Maryland streams is found in: 2000-2004 Maryland Biological Stream Survey Volume 14: Stressors (http://www/dnr/Maryland.gov/streams/pubs/ea05-11_biodiv.pdf).





Figure 8-4. Relative severity of stressors affecting biota in Maryland streams, based on 2000-2004 MBSS data




Figure 8-5. Stressors by county, in percent of total stream miles for that county, based on the 2000-2004 MBSS. Values will be greater than 100% because multiple stressors can impact the same stream reach